

1ST LECTURE:

MATHEMATICAL FOUNDATIONS AND INTERACTIVE EQUATIONS AND THEIR APPLICATIONS IN QUANTUM CHEMISTRY

الأسس الرياضية والمعادلات التفاعلية وتطبيقاتها في الكيمياء الكمية



Energy:

It is the ability to get a job done

There are types or states of energy, including:

1-kinetic energy

مسالح، الطانه الخركيه

2-Potential energy (vi) is the P.E P.E

Kinetic energy

at & jailed 1

Stat at 45therailed 1

m (kg) V (m/5)

KE= 1 mV2

(Usp) wrist op? $kg \cdot \frac{m^2}{5^2} = kg m^2/5^2 = J$

کیہ میا کیں

 $KE = \frac{1}{2}mV^2 = \frac{1}{2}(10)(20)^2$

= 2000+

potential energy

الغامة) س عدّلكها ,لم نعبه لوضعه (مختزنه) طامة الوضع) كلابيه الفامة) كروتيم

PEg=Ug=Eg

م طقه الوضع کازبه تعتمدی الکتکه م م م م م م م م م م

E=PE=mgh

م رفعناصم کن لاعل کتله و کالای ۱۵ مراک ۱۵ مراک مراک مراک کنونه فیم کالونه ک

E= mg h = 3(9.8)10 294 T

1-kinetic energy

عاق مبذلكها الحب سب الحك

kinetic energy, form of energy that an object or a particle has by reason of its motion

And from the laws of traditional (classical) mechanics, the kinetic energy of a body of mass (Kg) And it moves fast v (m/s) be ½ mv2

Kinetic Energy

K.E. = $\frac{1}{2}$ MV²

M: Mass (Kg)

V: Velocity (m/s)

- ☐ **kinetic energy,** form of energy that an object or a particle has by reason of its motion.
- Kinetic energy is a property of a moving object or particle and depends not only on its motion but also on its mass.
- ☐ The kind of motion may be translation (or motion along a path from one place to another), rotation about an axis, vibration, or any combination of motions.

خليط اعتزازه


Translational kinetic energy of a body is equal to one-half the product of its mass, m, and the square of its velocity, v, or 1/2mv2.

الطاقة الحركية: هي شكل من أشكال الطاقة التي يمتلكها جسم أو جسيم نتيجةً لحركته.

- الطاقة الحركية خاصية للجسم أو الجسيم المتحرك وتعتمد ليس فقط على حركته، بل أيضًا على كتلته.
- قد تكون الحركة على شكل: انتقال (حركة على مسار من مكان إلى آخر)، أو دوران حول محور، أو اهتزاز، أو مزيج من
 هذه الحركات.
 - الطاقة الحركية الانتقالية لجسم تساوي نصف حاصل ضرب كتلته m في مربع سرعته v، أي: •

$$E_k = \frac{1}{2}mv^2$$

 $E_k = kinetic\ energy\ of\ object$ $m = mass\ of\ object$ $v = speed\ of\ object$





The usual SI unit of kinetic energy, is the unit of mass (kg) multiplied by the unit of velocity م مر لعباس (m/sec) squared.

$$\log \left(\frac{\mathbf{m}}{\sec^2}\right)^2 = \frac{\log m^2}{\sec^2} = 3$$

Notice that this is the same as the unit of work, the Joule. All forms of energy can be measured in Joules.

Examples





1. What is the KE of a White-tailed Jackrabbit of mass 3.4 kg when running at a speed of 15 m/sec?


$$kE = \pm mv^{2}$$

$$= \frac{1}{2} (3.4 \text{ kg}) (15 \frac{m}{\text{sec}})^{2}$$

$$= \frac{1}{2} (3.4 \text{ kg}) 225 \frac{m^{2}}{\text{sec}^{2}}$$

$$= 382.5 \frac{\text{kg m}^{2}}{\text{sec}^{2}}$$

$$= 380 \text{ Tourded}$$



```
example

mass = 55 kg

velocity = 3.87m/s

= \frac{1}{2} (55)(3.87)^2

= \frac{1}{2} (55)(3.87)^2

= \frac{1}{2} (55)(3.87)^2
```

$$KE = 0.5 \times m v^2$$

= 0.5 × 55kg × (3.87m/s)²

$$KE = 0.5 \times \text{m v}^2$$

= 0.5 x 55kg x 14.97m²/s²
= 411.675 J



- Example 1: What is the <u>velocity</u> of an object with a mass of 30 kg and a <u>kinetic</u> energy of 500 J?
 - $KE = 0.5 \times mv^2$
 - $500 J = 0.5 \times 30 \times v^2$
 - Multiply mass by 0.5: 0.5 x 30 = 15
 - Divide kinetic energy by the product: 500/15 = 33.33
 - Square root to find velocity: 5.77 m/s
- Example 2: What is the mass of an object with a kinetic energy of 100 J and a velocity of 5 m/s?
 - $KE = 0.5 \times mv^2$
 - $100 J = 0.5 \times m \times 5^2$
 - Square the velocity: $5^2 = 25$
 - Multiply by 0.5: $0.5 \times 25 = 12.5$
 - Divide kinetic energy by product: 100/12.5 = 8 kg

example 1	
v ² = 500j	
0.5 x 30kg	
= <u>500j</u>	
15 kg	
$v2 = 33.33 \text{m}^2/\text{s}^2$	
$v = \sqrt{33.33} \text{m}^2/\text{s}^2$	
v= 5.77m/s	

example 1

$$m = 100j$$
 $0.5 \times (5 \text{ m/s})^2$
 $= 100j$
 $0.5 \times 25 \text{ m}^2/\text{s}^2$
 $= 100j$
 $12.5 \text{ m}^2/\text{s}^2$
 $= 8 \text{ kg}$

$$\frac{2KE - v^2}{m}$$

$$V = \int_{30}^{2 \times 500} = 5.773 \text{ m/s}$$

E+2

$$\frac{2KE}{V^2} = m$$

$$m = \frac{2 \, \text{KE}}{V^2} - \frac{2 \, \text{X} \, 100}{25} - 8 \, \text{Kg}$$



Types of Kinetic Energy

There are five types of kinetic energy that are,

- •Radiant Energy
- •Thermal Energy
- •Sound Energy
- •Electrical Energy
- Mechanical Energy

أنواع الطاقة الحركية

هناك خمسة أنواع رئيسية من الطاقة الحركية:

- 1. الطاقة الإشعاعية (Radiant Energy)
- 2. الطاقة الحرارية (Thermal Energy)
- 3. الطاقة الصوتية (Sound Energy)
- 4. الطاقة الكهربائية (Electrical Energy)
- 5. الطاقة الميكانيكية (Mechanical Energy)

Radiant Energy

Radiant energy is a form of kinetic energy, which is the physical energy of matter resulting from electromagnetic radiation. It is radiated from matter into the surrounding environment.

For example, Ultraviolet light, Gamma rays, and X-rays all have electromagnetic radiant energy.

Thermal Energy

Thermal energy can be also known as heat energy. It is generated by the kinetic energy of the motion of atoms when they collide with each other.

For Example, Hot Springs, Heated Swimming pools have thermal energy.

• هي شكل من أشكال الطاقة الحركية، وهي الطاقة الفيزيائية للمادة الناتجة عن الإشعاع الكهرومغناطيسي.

- تُشعّ من المادة إلى البيئة المحيطة.
- أمثلة: الضوء فوق البنفسجي، أشعة غاما، الأشعة السينية (X-rays).

2. الطاقة الحرارية

1. الطاقة الاشعاعية

- تُعرف أيضًا بطاقة الحرارة.
- تنتج من الطاقة الحركية لحركة الذرات عند تصادمها مع بعضها البعض.
 - أمثلة: الينابيع الساخنة، وأحواض السباحة المُسخنة.

Sound Energy

Sound energy is the energy, produced by the vibration of an object. It travels through the medium but cannot travel in a vacuum or space.

For Example, Tuning Forks, Beating Drums have sound energy.

Electrical Energy

Electrical energy can be obtained from the free electrons that are of positive and negative charge.

For Example, Batteries have electrical energy.

Mechanical Energy

The sum of kinetic energy and potential energy is called mechanical energy. It can neither be created

nor destroyed but it can be converted from one form to other.

For Example, Satellites orbiting around the Earth, Moving Cars all have mechanical energy.

3. الطاقة الصوتية (Sound Energy)

- هي الطاقة الناتجة عن اهتزاز جسم ما.
- تنتقل عبر وسط مادي (كالهواء أو الماء)، لكنها لا تستطيع الانتقال في الفراغ أو الفضاء.
 - أمثلة: الشوكة الرنانة، قرع الطبول.

4. الطاقة الكهربائية (Electrical Energy)

- 🕐 تُستمد من الإلكترونات الحرة ذات الشحنات الموجبة والسالبة.
 - أمثلة: البطاريات تمتلك طاقة كهربائية.

5. الطاقة الميكانيكية (Mechanical Energy)


- هي مجموع الطاقة الحركية والطاقة الكامنة.
- لا تُخلق ولا تُفنى، ولكن يمكن تحويلها من شكل إلى آخر.
- أمثلة: الأقمار الصناعية التي تدور حول الأرض، والسيارات المتحركة.

2-Potential energy

It is the energy resulting from the presence of the body in a specific position, and is often the result of forces The attraction or repulsion of things

Types of potential energy include:

- Gravitational potential energy
- Chemical energy
- Nuclear energy
- Elastic potential energy, also called spring energy



الطاقة الكامنة (Potential Energy)

- هي الطاقة الناتجة عن وجود الجسم في موضع معيّن،
- وغالبًا ما تكون نتيجة لقوى الجذب أو التنافر بين الأشياء.

أنواع الطاقة الكامنة تشمل:

- 1. الطاقة الكامنة الجاذبية (Gravitational Potential Energy):
 - ناتجة عن موقع الجسم في مجال الجاذبية.
 - مثال: كرة على ارتفاع معيّن عن سطح الأرض.
 - 2. الطاقة الكيميائية (Chemical Energy):
 - مخرّنة في الروابط الكيميائية بين الذرات والجزيئات.
 - مثال: الغذاء، الوقود.
 - 3. الطاقة النووية (Nuclear Energy):
 - ناتجة عن تفاعلات داخل نواة الذرة (انشطار أو اندماج).
 - مثال: محطات الطاقة النووية.
- الطاقة الكامنة المرنة (Elastic Potential Energy) أو طاقة النابض:
 ناتجة عن تمدد أو انضغاط جسم مرن مثل النابض أو الشريط المطاطى.
 - مثال: القوس المشدود، الزنبرك المضغوط.

Gravitational Potential Energy

Gravitational potential energy is the energy stored in an object due to its position above the Earth's surface.

$$PE = E_p = mgh$$


m = mass (kg)

m/52

g = gravitational field strength (N/kg)

h = height (m)

 E_p = gravitational potential energy (J)



Example 🚜

A book of mass $1~\mathrm{kg}$ is kept on a library shelf at height. If the change in potential energy is 17.64J. Then calculate the height of the bookshelf. We already know that the change in energy is equal to the potential energy of the object at that

height

$$\Delta E_{pe} = mgh$$

17. 64J = 1kg × 9. 8N/kg ×
$$h$$


$$h = \frac{17.64 \text{J}}{9.8 \text{N/kg}}$$


$$h = 1.8 \text{m}$$

The book is at a height of 1.8 m.

$$6in/8$$
, $5in/8$, $9in/8$, 9

$$h = 14.64 = 1.8 \text{ m}$$
 1×9.8





Example

Calculate the work done to raise an object of mass $5.5\,\mathrm{Kgto}$ a height of $2.0\mathrm{min}$ the earth's gravitational field.

We know that the work done to raise an object to a certain height is the gravitational potential energy of the object at that height so

Mass =
$$5.50 \,\mathrm{kg}$$

Height =
$$2.0m$$

$$g = 9.8 \text{N/kg}$$

$$work = \Delta E = mgh = 5.5 \times 9.8 \times 2$$

$$= 1000$$

Substitute these values in the equation for potential energy and we get

$$E_{\mathrm{pe}}\!=\!mgh$$

$$E_{\mathrm{pe}}\!=\!5.50kg imes9.8\mathrm{N}/kg imes2.0\mathrm{m}$$

$$E_{\mathrm{pe}}\!=\!110\mathrm{J}$$


Therefore the work done to raise an object of mass $5.5\,\mathrm{kgto}$ a height of $2\mathrm{mis}$ $110\mathrm{J}$.


O Di Th

Difference Between Kinetic Energy and Potential Energy

The difference between Kinetic Energy and Potential Energy is discussed below,

Kinetic Energy	Potential Energy
Kinetic energy is the energy of any moving object.	Potential energy is the energy of the object due to its position in space.
Formula for kinetic energy is, K.E. = 1/2mv ²	Formula for potential energy is, P.E. = mgh
Example: Energy of a moving car.	Example: Energy of a book placed on top of the table.





➤ What is potential energy?

Potential energy E_P, is a form of energy that depends on the relative position of an object within a system.

➤ What is an example of potential?

ما هي الطاقة الكامنة؟

Examples of potential energy are

الطاقة الكامنة $_{P}E$ هي شكل من أشكال الطاقة التي تعتمد على الموضع النسبي لجسم داخل نظام.

Raised object
Stretched rubber band
Water stored in a dam
The energy released during nuclear fusion and fission of atoms

ما هي أمثلة على الطاقة الكامنة؟

جسم مرفوع عن سطح الأرض.

شريط مطاطي مشدود.

المياه المخزنة في السدود.

الطاقة المنطلقة أثناء الاندماج أو الانشطار النووي للذرات.

➤ What is the formula for calculating potential energy?

The potential energy can be calculated by EP = mgh

ما هي الأنواع الأربعة للطاقة الكامنة؟

الأنواع الأربعة للطاقة الكامنة هي:

- 1. الطاقة الكامنة الجاذبية (Gravitational Potential Energy)
 - 2. الطاقة الكامنة المرنة (Elastic Potential Energy)
 - 3. الطاقة الكامنة الكهربائية (Electric Potential Energy)
 - 4. الطاقة الكامنة النووية (Nuclear Potential Energy)

- ➤ What are the 4 types of potential energy?
- The 4 types of potential energy are

Gravitational Potential Energy
Elastic Potential Energy
Electric Potential Energy
Nuclear Potential Energy

ما الفرق بين الطاقة الكامنة والطاقة الحركية؟

- الطاقة الكامنة: هي طاقة مخزنة ناتجة عن الوضع النسبي لجسم داخل نظام.
 - الطاقة الحركية: هي طاقة مرتبطة بحركة الجسم.

➤ What is the difference between potential and kinetic energy?

Potential energy is a stored form of energy due to the relative position of an object within a system whereas, kinetic energy is due to the motion of the object



☐ In the International System of Units SI joules (J)

$$1J = kg.m^2.s^{-2}$$

$$1J = (kg.m.s^{-2}).m$$

$$1J=N.m$$

In quantum chemistry, we usually use the electron volt (eV)

Power (energy) abl $= kgm^2 = kgm^2 s^{-2}$ J= Nom 1.602 × 10 1 lev 1.602 X 10-19 J 60 = 4.184 J 1 cal Cal 4.184

الصانة الله يعيلها المحتور: عنه حركة نهائي عهم حقواره 110

An electron volt : is the energy obtained by an electron when it passes under the influence of a voltage of one volt (1V).

 $1 \text{ eV} = 1.602 \text{ x } 10^{-19} \text{ coulomb x } 1\text{V}$ = $1.602 \text{ x } 10^{-19} \text{ CV} = 1.602 \text{ x } 10^{-19} \text{ J}$

J = eV = polog . Fri


☐ Another commonly used unit is calorie (cal).

المان المازم لرفي دام عراء على المان المازم المازم

It is defined as the amount of energy needed to increase the temperature of one gram of water by one degree Celsius

1 cal = 4.184 J

Units of Energy



- 1. ERG a unit of energy (the smallest unit of energy)
- 2. ELECTRON VOLT (eV) $1 \text{ eV} = 1.60 \times 10^{-12} \text{ ergs}$

(the eV is smaller than the erg, but it is a relative unit whose value must be determined experimentally.)

3. CALORIE - the most commonly used unit of energy.

$$1 \text{ Cal} = 4.18 \times 10^7 \text{ ergs}$$

4. JOULE - 1 Joule = 1.00×10^7 ergs

4.18 Joules = 1 Calorie

Waves and Particles

In classical physics, the concepts of waves and particles are mutually exclusive.

Waves Waves

Wavelength Frequency

• Spread in space and time.

Pass through each other

• Can be superposed – show interference effects.

O Particles

Localized in space and time.

Cannot pass through each other - they bounce or shatter.

In classical physics, nothing can be both a particle and a wave at the same time.

المحم محلاسته

Properties of waves

The five important properties of waves are Amplitude, Frequency, Wavelength, period, and Speed. Listed below is a detailed view of all the properties.

Amplitude Amplitude is among the first properties of waves and is denoted by the symbol (A.).

Hs S.I. unit is a meter (m)

2\Frequency

The second property of waves is the frequency denoted by the symbol of The Greek symbol is lambda valled nu.

- •Its S.I. unit is Hertz (Hz), named in honour of a German Physicist Heinrich Rudolf Hertz.
- •It measures the total number of waves passing through a definite point in a given period.
- •Frequency is **inversely** proportional to the period and is given by the formula f=1/T.
- •The relation between wavelength, speed, and frequency can be determined by λ .
- In the case of in electromagnetic wave that propagates via vacuum, the sis replaced by c (speed of light), and the formula changes to $f=c/\lambda$.

≺ خصائص الموجات

الخصائص الخمس المهمة للموجات هي: السعة (Amplitude)، التردد (Frequency)، الطول الموجي (Wavelength)، الزمن الخصائص الخمس المهمة للموجات هي: السعة (Speed)، والسرعة (Speed)، فيما يلى عرض تفصيلي لكل خاصية:

- 1. السعة (Amplitude)
- السعة من أولى خصائص الموجات، ويرمز لها بالرمز A.
- وحدتها في النظام الدولي للوحدات (SI) هي المتر (m).
 - 2. التردد (Frequency)
- الخاصية الثانية من خصائص الموجات هي التردد، ويرمز له بالرمز f.
 - الرمز الإغريقي المستخدم هو **v** (نيو nu).
- وحدته في النظام الدولي للوحدات (SI) هي الهرتز (Hz)، سُمّيت تكريمًا للفيزيائي الألماني هاينريش رودولف هرتز.
 - · ا<mark>ت</mark>ردد يقيس عدد الموجات الكلي التي تمر عبر نقطة معينة خلال فترة زمنية محددة.
 - التردد يتناسب عكسيًا مع الزمن الدوري، ويُعطى بالعلاقة: f = 1/T.
 - العلاقة بين الطول الموجي والسرعة والتردد تُعطى بالصيغة: f = v/λ.
- في حالة الموجات الكهرومغناطيسية التي تنتشر في الفراغ، تُستبدل ν بـ c (سرعة الضوء)، وتصبح الصيغة: f = c/λ.

3.Wavelength

The wavelength is the third property of waves. It is the total displacement or distance between two corresponding troughs or crests.

- •It is denoted by the Greek symbol lambda (λ)
- •Its S.I. unit is a meter (m)

4. Time period

- The time period is the fourth property of waves and is defined as the time a particle of a medium takes to make a whole vibrational cycle.
- •Its S.I unit is seconds (s)
- •The frequency and time period are **inversely** related, T=1/f; if frequency increases, the time period decreases, T represents the time period, and f denotes frequency.

5.Speed/Velocity

- الطول الموجي هو الخاصية الثالثة للموجات. وهو المسافة الكلية بين قمتين متتاليتين أو قاعين متتاليين للموجة.
- Speed id, the fifth property of a wave, is defined as the total distance covered by a point on the wave in a specific period of time.

وحدته في النظام الدولي للوحدات (SI) هي المتر (m).

3. الطول الموجى (Wavelength)

پُرمز له بالرمز الإغريقي λ (لامبدا).

4. الزمن الدوري (Time Period)

•The S.I. unit of speed is metres per second (m/s).

- الزمن الدوري هو الخاصية الرابعة للموجات، ويُعرَّف بأنه الزمن الذي يستغرقه جسيم من الوسط لإكمال دورة اهتزازية كاملة.
- •The formula for calculating the speed of a wave is speed = Wavelength frequency.

•It depends on the medium through which the wave is propagating.

- وحدته في النظام الدولي للوحدات (SI) هي الثواني (s).
 - العلاقة بين التردد والزمن الدوري عكسية: T = 1/f.
- إذا زاد التردد قل الزمن الدوري. حيث T يمثل الزمن الدوري و f يمثل التردد.
 - 5. السرعة/السرعة الموجية (Speed / Velocity)
- السرعة هي الخاصية الخامسة للموجات، وتُعرَّف بأنها المسافة الكلية التي يقطعها نقطة على الموجة خلال فترة زمنية
 - وحدتها في النظام الدولي للوحدات (SI) هي متر لكل ثانية (m/s).
 - تعتمد على الوسط الذي تنتشر خلاله الموجة.
 - الصيغة الرياضية لحساب سرعة الموجة هي:
 - السرعة = الطول الموجى × التردد

Properties of Waves


A wave can be thought of as a vibrating disturbance by which energy is transmitted.


Wave characteristics

Wavelength λ (lambda) is the distance between identical points on successive waves.

Frequency v (nu) is the number of waves that pass through a particular point in one second.

Amplitude is the vertical distance from the midline of a wave to the peak or trough.





Making the point of the same of the same

E = h \mathcal{V} $h = 6.626 \times 10^{-34} \text{ t.s}$

 $E = hv = hc = hc\bar{v}$


The speed of a wave depends on the type of wave and the nature of the medium through which the wave is traveling (for example, air, water, or a vacuum).

The **speed** (u) of a wave is the product of its wavelength and its frequency:


$$u = \lambda v$$

Wavelength is usually expressed in units of meters, centimeters, or nanometers, and frequency is measured in hertz (Hz), where

$$1 \text{ Hz} = 1 \text{ cycle/s}$$



Two waves having different wavelengths and frequencies. The wavelength of the top wave is three times that of the lower wave, but its frequency is only one-third that of the lower wave. Both waves have the same speed and amplitude.



Wave number: is the reciprocal of the wavelength and is denoted by the symbol



Its units are the reciprocal of units of length and is defined as the number of waves in 1 meter

Energy: It is calculated from the relationship

$$E = hv$$

$$h = 6.626 \times 10^{-34} J \cdot s$$

Where h : Planck's constant is equal to

The relationship between the previous properties

$$E = hv = \frac{hc}{\lambda} = hc\overline{v}$$

a)
$$E = \frac{LC}{D} = \frac{6.626 \times 10^{-34} \times 3 \times 10^{8}}{556 \times 10^{-9}} = 3.164 \times 10^{-19}$$

2-550X10 m

C = 3 x 108 m/s

Application:

Application:
b)
$$v = \frac{3 \times 10^{3}}{5} = \frac{3 \times 10^{3}}{550 \times 10^{3}} = 5.45 \times 10^{4}$$
 Hz
A sodium lamp emits a yellow light of wavelength 550 nm?

Calculate E , ν and $\overline{\nu}$

c)
$$\sqrt{1} = \frac{1}{1} = \frac{1}{550 \times 10^{-9}} = 1.812 \times 10^{-9}$$
 $m-1$

☐ Convert wavelength to metre

$$\lambda = 550 \text{ nm} = 550 \text{ x } 10^{-9} \text{ m} = 5.5 \text{ x } 10^{-7} \text{ m}$$

 $1 \text{ nm} = 10^{-9} \text{ m}$

1- the wave number (equal to the reciprocal of the wavelength)

$$\overline{v} = \frac{1}{\lambda} = \frac{1}{5.5 \times 10^{-7} \,\mathrm{m}} = 1.82 \times 10^6 \,\mathrm{m}^{-1}$$




2-Frequency:

$$v = \frac{c}{\lambda} = c\overline{v} = 3 \times 10^8 \,\text{ms}^{-1} \times 1.82 \times 10^6 \,\text{m}^{-1} = 5.45 \times 10^{14} \,\text{s}^{-1}$$

3- energy:

$$E = hv = 6.626 \times 10^{-34} Js \times 5.45 \times 10^{14} s^{-1} = 3.61 \times 10^{-19} J$$






What is the frequency?

Frequency is defined as the number of oscillations of a wave per unit of time and is measured in hertz(Hz)


What is a wavelength?

Wavelength is defined as the distance between two nearest points in phase with each other.

What is the relationship between frequency and wavelength? Inverse relationship between frequency and wavelength? The wave speed is equal to the product of its frequency and wavelength. $V = f \lambda$







A radio station broadcasts at a frequency of 99,500,000Hz. If the broadcast is an electromagnetic wave, then what is its wavelength?

$$c=3*10^8\frac{m}{s}$$

$$\lambda = \frac{C}{f} = \frac{3 \times 10^8}{99500000} = 3.02 \text{ m}$$

Explanation:


The relationship between wavelength and frequency is given by the equation $\lambda=\frac{c}{f}$, where λ is the wavelength, c is the speed of light, and f is frequency.

We are given the values for frequency and the speed of light, allowing us to solve for the wavelength.

$$\lambda = \frac{c}{f}$$

$$\lambda = \frac{3 * 10^8 \frac{m}{s}}{99,500,000 Hz}$$

$$\lambda = 3.02m$$





A wave oscillates with a speed of $8.9\frac{m}{s}$ and has a wavelength of 6.7m. What is the frequency of the wave?

$$f = \frac{U}{\lambda} = \frac{8.9}{6.7} = 1.33 \text{ HZ}$$

Explanation:

The equation for velocity in terms of wavelength and frequency is $v=\lambda f$.

We are given the velocity and the wavelength. Using these values, we can solve for the frequency.

$$v = \lambda f$$

$$8.9\frac{m}{s} = 6.7m * f$$

$$\frac{8.9\frac{m}{s}}{6.7m} = f$$

$$1.33Hz = f$$

e|x|e|r|c|i|s|e

Example

A sound wave has a time period of 0.0001 seconds. What is its frequency?

Answer


$$f=\tfrac{1}{T}$$

$$f = \frac{1}{0.0001 s}$$


$$f = 10,000 Hz$$


The frequency of the sound wave is 10,000 Hz.

$$\hat{f} = \frac{1}{T} = \frac{1}{0.0001}$$



Sketch a diagram of a wave and label the amplitude and wavelength.





Q1. What is the relationship between frequency and wavelength? (Direct or Inverse)

Q2. What is the relationship between frequency and energy? (Direct or Inverse)

What Is Light?

There was a long debate about the nature of light This debate over the particle or wave nature of light came to a resolution in 1803 when the experiment confirmed that light propagates in waves

≺ ما هو الضوء؟

كان هناك جدل طويل حول طبيعة الضوء، هل هو ذو طبيعة جسيمية أم موجية. وقد حُسم هذا الجدل في عام 1803 عندما أثبتت التجارب أن الضوء ينتشر على شكل موجات.

1- الطبيعة الموجية للضوء

الإشعاع الكهرومغناطيسي

1-The Wave Nature of Light

الإشعاع الكهرومغناطيسي هو أحد أشكال الطاقة.
 ووفقًا للنظرية الكهرومغناطيسية التقليدية، فإن الطاقة الكهرومغناطيسية تُحمَل بواسطة موجات.

وتنقل هذه الموجات الطاقة في اتجاه انتشارها.


Electromagnetic Radiation


Electromagnetic radiation is one of the forms of energy, and based on the traditional electromagnetic theory, electromagnetic energy is carried by waves and transfers energy in the direction of wave propagation

اعدمات المحدمفامية سرّ ل في لراني


• Electromagnetic waves travel through a vacuum at a constant speed called the speed of light, c which has the defined value of exactly $2.99792458 \times 10^8 \text{ m s}^{-1}$.


C= 2-0979 x108 m/s






Note the small fraction that is visible to the human eye. The electromagnetic radiation that is visible to the human eye has a wavelength in the range 380 nm (violet light) to 750 nm (red light).





Which of the following wave has minimum energy?

- 1.) Radio waves
 - 2. UV rays
 - 3. Infra-red



What is the relationship between frequency and wavelength?

Frequency and wavelength are inversely related. If the wavelength is longer, the frequency is less. A shorter wavelength has a greater frequency.

What is the relationship between frequency and energy?

They are proportionally related. As one becomes greater, so does the other. Energy of a wave is measured by its frequency.


At is the relationship between wavelength and energy?


What is the relationship between wavelength and energy?

They are inversely related. Because energy is measured using frequency, and wavelength is inversely related to frequency; this means that wavelength and energy are also inversely related. As wavelength gets longer, there is less energy.

العلاقة	الصيغة/الوصف	النتيجة
التردد ↔ الطول الموجي	ass has	كلما زاد الطول الموجي → قل التردد، وكلما قصر الطول الموجي → زاد التردد
التردد ↔ الطاقة	as y true	كلما زاد التردد → زادت الطاقة، وكلما قل التردد → قلت الطاقة
الطول الموجي ↔ الطاقة	SAL ME	كلما زاد الطول الموجي → ق <mark>لت</mark> الطاقة، وكلما قصر الطول الموجي → زادت الطاقة


ercise





Label the parts of a wave using the terms: crest, trough, frequency, wavelength, amplitude

- 1. wavelength
- 2. trough
- amplitude
- 4. crest
- 5. frequency



2-Particle Nature of Light

Classical Physics

In the first half of the 19th century, many discoveries ware made such as:

- Three laws of motion (mechanics) by Newton.
- Three laws of thermodynamics by Rumfold, Joule, Gibbs and Carnot.
- Kinetic theory of gases and statistical mechanics by Maxwell, Gibbs and Sylveline Ezis Boltzmann.
- Wave theory of light by Huygen.
- Optics by Young and Frensel.


The body of these accomplishme what we now call classical phys

2- الطبيعة الحسيمية للضوء

الفيزياء الكلاسيكية

- في النصف الأول من القرن التاسع عشر، تم القيام بالعديد من الاكتشافات مثل:
 - قوانين الحركة الثلاثة (الميكانيكا) بواسطة نيوتن.
 - القوانين الثلاثة للديناميكا الحرارية بواسطة رمفورد، جول، جيبس، وكارنو.
- النظرية الحركية للغازات والميكانيكا الإحصائية بواسطة ماكسويل، جيبس، وبولتزمان.
 - النظرية الموجية للضوء بواسطة هويجنز.
 - علم البصريات بواسطة يونغ وفريسنل.
- تُعتبر مجموعة هذه الإنجازات في الفيزياء هي الأساس لتطور ما نُسميه الآن بـ الفيزياء الكلاسيكية.

- By the turn of the 19th century it was pretty much established that **light was an electromagnetic wave**.
- However, three crucial experiments showed that this description of light was incomplete.
- This first was Blackbody Radiation (thermal radiation of hot objects).
- The second was the Photoelectric Effect.
- The third one was the Compton Effect.



- مع بدایة القرن التاسع عشر، کان قد تم إثبات أن الضوء موجة کهرومغناطیسیة بشکل کبیر.
 - لكن، ثلاث تجارب أساسية أظهرت أن هذا الوصف للضوء غير مكتمل:
 - 1. إشعاع الجسم الأسود (الإشعاع الحراري للأجسام الساخنة).
 - 2. التأثير الكهروضوئي.
 - 3. تأثير كومبتون.

Important Mathematical Foundations in Quantum Chemistry

In our study of quantum chemistry we will encounter some related terms Certain meanings, the most important of these terms are:

1 - differential

2 - Integration

3 - Complex numbers

4 - Vector 5 - Operators

6 - conservative functions

7 - odd and even functions

8 - Reference systems

الأسس الرياضية المهمة في الكيمياء الكمية

في دراستنا للكيمياء الكمية سنواجه بعض المصطلحات ذات الصلة،

ومن أهم هذه المصطلحات:

- 1. التفاضل
- 2. التكامل
- 3. الأعداد المركبة
 - 4. المتجهات
- 5. المؤثرات (Operators)
 - 6. الدوال المحافظة
- 7. الدوال الفردية والزوجية
- أنظمة الإحداثيات (أنظمة المرجع)

First: the differential

differential rule

$$f(n) = y = x^n$$

differential Xⁿ

$$y' = dy - f(x)$$

nx ^{n-l}

$$f(x) = n x^{n-1}$$

 $f(x) = x^2 + 3x^4 + 5$
 $f'(x) = 2x + 3$

$$f(x) = 3x^5$$

 $f'(x) = 15x^4$

Some important differential laws (note that a is a constant)

Sinx cosh
$$cosh - sinx$$

$$cosh - sinx$$

$$d = 30x^{2}$$
Sin4h 4cos4x
$$d = 30x^{3} = 30x^{2}$$

$$cosh - sinx$$

$$d = 30x^{2}$$

$$d = 30x^{$$

$$\frac{d}{dx}a = 0$$

$$\frac{d}{dx}(f.9) = fg' + f'g$$

$$\frac{d}{dx}ax^3 = 3ax^2$$

$$\frac{d}{dx}\sin ax = a\cos ax \frac{dx}{dx}(x^2e^{-\alpha x^2})$$

$$\frac{d}{dx}e^{-ax^2} = -2ax e^{-ax^2}$$

$$\frac{d}{dx}e^{-ax^2} = x^2(-2axe^{-ax^2}) + e^{-ax^2}(2x)$$

$$= -2ax^3e^{-ax^2} + 2xe^{-ax^2}$$

$$=\int x^n dx = \frac{x^{n+1}}{x^{n+1}}$$

integration rule

The integration of
$$X^n = \int x^3 dx = \frac{x^4 + C}{4}$$

$$\frac{x^{n+1}}{n+1} \int \frac{1}{x^3} dx = \int x^{-3} dx = \frac{x^{-2}}{-2}$$

☐ This rule is true for both negative and positive n values

$$\int \frac{1}{X} dx = lnx + c$$

$$\int \frac{f'}{f} dx = \int \frac{2x}{x^2} dx = 2x^2 + c$$

Second: integration $\int 4 dx = 4x$ $\int 4 dy = 4y$

Some important laws of integration (note that

$$\int_{0}^{3x} = \frac{3x}{3}$$

a is a constant)
$$\int Sin x \, dx = -\cos x \, t \, C$$

$$\int C \times dx = e^{x}$$

$$\int C \cos x \, dx = \sin x + C$$

$$\int a dx = 5x + C$$

$$\int ax^3 dx = 0 + C$$

$$\int \sin ax dx = -\cos x + C$$

$$\int \cos ax dx = -\sin x + C$$

$$\int adx = ax + c$$

$$\int ax^{3}dx = a\frac{x^{4}}{4} = \frac{a}{4}x^{4} + \zeta$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax + C$$

$$\int \cos ax \, dx = \frac{1}{a} \sin ax + C$$

types of integration

Julimited integration:

The limits of integration are not stated

b)limited integration:

The limits of integration are indicated

The limits of integration are indicated

$$x^n e^{-ax} dx = \frac{n!}{a^{n+1}}, n - 1, a = 0$$
 $\int_{0}^{\infty} x^n e^{-ax} dx = \frac{e^{ax}}{a}$

$$\int_{3}^{3} 3 \times^{2} d \times = /3 \times 3$$

$$2 \times 3$$

$$2 \times 3$$

$$21 - 8 - 10$$

$$\int xe^{ax} dx = \frac{e^{ax}}{a} \left(x - \frac{1}{a} \right)$$


Application


Find the product of the following integrals.

$$\int_{1}^{4} x^{2} dx = \frac{x^{3}}{3} \int_{1}^{4} = \frac{4^{3}}{3} - \frac{1^{3}}{3} = \frac{63}{3} = 21$$

$$1650 (e^{2}-1) \int_{0}^{10} x dx = \frac{x^{2}}{2} \int_{0}^{10} = \frac{10^{2}}{2} - \frac{0^{2}}{2} = 50$$


$$\int_{0}^{2} e^{x} dx = e^{x} \int_{0}^{2} - e^{2} e^{2} = e^{2} - 1$$





Label the parts of a wave using the terms: crest, trough, frequency, wavelength, amplitude

- 1 wavelength
- 2. trough
- 3. amplitude
- 4. crest
- 5. <u>frequency</u>



2-Particle Nature of Light

Classical Physics

In the first half of the 19th century, many discoveries ware made such as:

- Three laws of motion (mechanics) by Newton.
- Three laws of thermodynamics by Rumfold, Joule, Gibbs and Carnot.
- Kinetic theory of gases and statistical mechanics by Maxwell, Gibbs and Boltzmann.
- Wave theory of light by Huygen.
- Optics by Young and Frensel.

The body of these accomplishments in physics is considered the development of what we now call **classical physics**.

- By the turn of the 19th century it was pretty much established that light was an electromagnetic wave.
- However, three crucial experiments showed that this description of light was incomplete.
- This first was Blackbody Radiation (thermal radiation of hot objects).
- The second was the Photoelectric Effect.
- The third one was the Compton Effect.

Important Mathematical Foundations in Quantum Chemistry

In our study of quantum chemistry we will encounter some related terms Certain meanings, the most important of these terms are:

1 - differential 2 - Integration 3 - Complex numbers


4 - Vector

5 - Operators

6 - conservative functions

7 - odd and even functions

8 - Reference systems



First: the differential

differential rule

differential Xⁿ

 nx^{n-l}

Some important differential laws (note that a is a constant)

$$\frac{d}{dx}a$$

$$\frac{d}{dx}ax^{3}$$

$$\frac{d}{dx}\sin ax$$

$$\frac{d}{dx}e^{-ax^{2}}$$

$$\frac{d}{dx}e^{-ax^{2}}$$

$$\frac{d}{dx}x^{2}.e^{-ax^{2}}$$

$$\frac{d}{dx}a = 0$$

$$\frac{d}{dx}ax^3 = 3ax^2$$

$$\frac{d}{dx}\sin ax = a\cos ax$$

$$\frac{d}{dx}e^{-ax^2} = -2ax e^{-ax^2}$$

$$\frac{d}{dx}x^{2}.e^{-ax^{2}} = x^{2}(-2axe^{-ax^{2}}) + e^{-ax^{2}}(2x)$$

$$= -2ax^{3}e^{-ax^{2}} + 2xe^{-ax^{2}}$$


Second: integration

integration rule

The integration of Xⁿ

$$\frac{x^{n+1}}{n+1}$$

- ☐ This rule is true for both negative and positive n values
 - \square Except n=-1



Second: integration

Some important laws of integration (note that a is a constant)

```
\int a dx
\int ax^3 dx
\int \sin ax dx
\int \cos ax dx
```

$\int adx = ax$

$$\int ax^{3}dx = a\frac{x^{4}}{4} = \frac{a}{4}x^{4}$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax$$

$$\int \cos ax \, dx = \frac{1}{a} \sin ax$$

types of integration

a)Unlimited integration:

The limits of integration are not stated

b)limited integration:

The limits of integration are indicated

$$\int_{0}^{\infty} x^{n} e^{-ax} dx = \frac{n!}{a^{n+1}}, n \rangle -1, a \rangle 0 \qquad \int x e^{ax} dx = \frac{e^{ax}}{a} \left(x - \frac{1}{a} \right)$$

Application

Find the product of the following integrals.

$$\int_{1}^{4} x^{2} dx$$

$$\int_{0}^{10} x dx$$

$$\int_{0}^{2} e^{X} dx$$